

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 7/50, 7/075, C11D 17/00, 9/26, 9/44		A1	(11) International Publication Number: WO 95/03782
			(43) International Publication Date: 9 February 1995 (09.02.95)
(21) International Application Number: PCT/AU94/00421			(81) Designated States: AU, CA, GB, KR, NZ, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(22) International Filing Date: 26 July 1994 (26.07.94)			
(30) Priority Data: PM 0246 28 July 1993 (28.07.93) AU			
(71) Applicant (for all designated States except US): ICI AUSTRALIA OPERATIONS PROPRIETARY LIMITED [AU/AU]; 1 Nicholson Street, Melbourne, VIC 3000 (AU).			
(72) Inventors; and (75) Inventors/Applicants (for US only): WANSBOROUGH, Katherine, Margaret [AU/AU]; 249 Forest Road, Boronia, VIC 3155 (AU). LOMBARD, Anne, Maree [AU/AU]; 23 Rowell Street, Rosanna, VIC 3084 (AU). JONES, Pamela, Joyce [AU/AU]; 12 Nartee Place, Wilberforce, NSW 2756 (AU).			
(74) Agent: DAVY, John, R.; Industrial Property Division, ICI Australia Operations Proprietary Limited, 1 Nicholson Street, P.O. Box 4311, Melbourne, VIC 3001 (AU).			
(54) Title: PEARLESCENT BASED CONCENTRATE FOR PERSONAL CARE PRODUCTS			
(57) Abstract A pearlescent based concentrate and method of preparation wherein the concentrate comprises at least one alkylpolysaccharide, traditional pearlescent ester or acid and alkyl sulphates. The concentrate is suitable for use in personal care products of low irritancy to the skin of users.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

PEARLESCENT BASED CONCENTRATE
FOR PERSONAL CARE PRODUCTS

The invention relates to pearlescent based concentrates comprising at least one alkylpolysaccharide, traditional
5 pearlescent esters or acid and alkyl sulphates and to processes for the preparation thereof.

Personal care products such as handsoaps, bubble baths, shampoos and hair conditioners often include in their composition agents which impart a pleasant, pearl-like glossy
10 appearance in order to enhance the appeal of such products to the consumer. The agents which impart this appearance are known as pearlescent based concentrates and generally comprise microscopic platelike crystals which remain
15 dispersed in compositions to which they are added and reflect light in such a manner as to impart a pearl-like appearance.

Pearlescent based concentrates of the prior art have included finely divided natural substances or inorganic compounds such as mica, fish scale, bismuth oxychloride or the like. For example, Japanese patent application no.
20 90/134,825 (Priority Date; 24 May 1990) describes compositions for cosmetics having a pearly appearance, comprising glucose fatty acid esters and mica and/or mica Ti.

In recent years, the use of a fatty acid glycolic ester in pearlescent concentrates has been particularly favoured.
25 In general such concentrates are produced by adding a fatty acid glycolic ester which is solid at room temperature to a composition such as shampoo or the like, followed by heating of the mixture to melt the fatty acid glycolic ester, and then by cooling to recrystallise it and thereby providing
30 small crystals which impart a pearl-like gloss to the composition. In an alternative method of production, a pearling agent dispersion (such as a fatty acid glycolic ester dispersion, which is dissolved and cooled to recrystallise in advance) is mixed with a shampoo or the like
35 at room temperature as described in Japanese Patent Application nos. 71021/81 and 216728/83.

Other known pearlescent based concentrates have comprised microcrystalline polyvalent metal salts of higher fatty acids or fatty acid glycolic esters such as those described in United States Patent 4,959,200 (Date of Patent; 25 September 1990).

Other commonly used pearlescent based concentrates have been prepared using ethylene glycol monostearate, ethylene glycol di-stearate, glycerol monostearate, glycerol di-stearate or a cetyl stearyl alcohol or the like in conjunction with other similar esters. These pearlescent based concentrates of the prior art have also normally comprised ether sulphates, betaines, ethanolamide, amine based surfactants and the like which can be used as emulsifiers. United States Patent 5,017,305 (Date of Patent; 21 May 1991) describes a typical pearlescent concentrate in the form of a free-flowing dispersion comprising pearlescing esters, ether carboxylic acids and one or more monoethanolamides of C_{12} to C_{22} fatty acids.

One of the disadvantages of the pearlescent based concentrates of the prior art is that they often comprise chemicals which are potential skin sensitisers. Japanese Patent Application no. 21678/83 for example, discloses a pearling agent containing a salt of alkyl sulphate or a salt of polyoxyalkylene alkyl sulphate, a fatty acid dialkanolamide and water as solvents, and a fatty acid glycolic ester at a high concentration. Many factors influence the irritation effect of surfactants and the molecular structure of the surface active agent can be closely linked to its effect on adsorption, solubilization, penetration, swelling, denaturation and general irritation on human skin.

Generally, the order of increasing human skin irritation of anionic surfactants commonly used in personal care products is in the following order:


Ammonium Laureth Sulphate < Sodium Laureth Sulphate < Ammonium Lauryl Sulphate < Sodium Lauryl Sulphate.

Furthermore, some of the pearlescent based concentrates of the prior art have included surfactants which may contain toxic trace impurities which are suspected carcinogens including nitrosamine containing compounds such as diethanolamides. European Patent no. 221465 (22 October 1992) for example, relates to a process for the production of pearlescent pigment comprising ethanolamide.

In general, alkyl polyglucosides demonstrate very little tendency to scalp irritations.


It has now been found that low irritant pearlescent based concentrates containing biodegradable surfactants can be prepared using alkylpolysaccharides. Ultimate biodegradability was measured using OECD tests 301E and 301A, 1981. Alkylpolysaccharides are particularly attractive for use in personal care products because they contain no toxic trace impurities and because they are derived from renewable resources such as coconut oil and wheat starch.

The current invention therefore provides a pearlescent based concentrate including:
between 5 and 40% by weight of the composition of alkylpolysaccharide of formula



where R^1 is selected from the group comprising linear or branched C_6 to C_{22} alkyl or alkenyl group
where G is selected from the group comprising a C_5 or C_6 saccharide and
 x is from 1 to 10;

between 3 and 30% by weight of the composition of alkyl sulphate and/or alkyl ether sulphate;
and between 5 and 20% by weight of pearlescing esters of formula



where R^2 is chosen from the group including C_{16} to C_{22} fatty acyl groups,
 R^3 is chosen from the group comprising H or R^2 ,

n is 2 or 3 and
x is from 1 to 120.

The current invention further provides a pearlescent based concentrate including:

- 5 $H-G_x-OR^1$
where R^1 is selected from the group comprising linear or branched C_6 to C_{22} alkyl or alkenyl group
where G is selected from the group comprising a C_5 or C_6 saccharide and
10 x is from 1 to 10;
between 3 and 30% by weight of the composition of alkyl sulphate and/or alkyl ether sulphate;
and between 5 to 20% by weight of pearlescing acid of formula
R COOH
15 where R is chosen from the group comprising C_{16} to C_{22} alkyl groups.

In a preferred embodiment the pearlescing esters or acid of the current invention is chosen from ethylene glycol monostearate, ethylene glycol distearate, stearic acid and
20 mixtures thereof.

In a preferred embodiment the pearlescent based concentrate of the current invention further includes secondary alkane sulphonates of the formula


- $R^4 - CH(CH_3)SO_3M$
25 where R^4 is chosen from the group comprising
 C_6 to C_{22} alkyl groups
and M is chosen from the group comprising alkali metals

In a further preferred embodiment the fatty alcohol ethoxylates used in pearlescent based concentrate of the
30 current invention are of the formula

- $R^5 (OC_2H_4)_y OH$
where R^5 is chosen from the group including
 C_6 and C_{22} alkyl or alkenyl groups
and y is from 1 to 120

The pearlescent based concentrate of the current invention may further include between 0 and 30% by weight of the composition of one or more optional additives chosen from the group including:

- 5 alkylsulphosuccinate or polyoxyalkylenealkylsulphosuccinate of the formula




where R^6 is chosen from the group including C_8 to C_{20} linear or branched alkyl groups,

- 10 R^7 is H or CH_3 ,

M^1 and M are independently chosen from the group comprising alkali metals, alkaline earth metals, NH_4^+ , C_1 to C_3 alkyl groups, substituted ammonia and hydroxy substituted C_2 or C_3 alkyl substituted ammonia and

- 15 n is from 0 to 8;

sorbitan fatty acid esters of formula,



where R^8 is chosen from the group including C_{12} to C_{18} ;

- secondary alkane sulphonates; fatty alcohol alkoxylates;
20 fatty acid alkoxylates; fatty alcohols; alkanoyl N-methylglucamides; ether carboxylic acids; ethoxylated sorbitan fatty acid esters; and alpha-olefin sulphonates.


In a preferred embodiment the optional additives of the pearlescent concentrate of the current invention includes

- 25 fatty alcohols of the formula




where R^9 is chosen from the group comprising C_6 to C_{22} alkyl groups

- In a further preferred embodiment the optional additives
30 of the pearlescent concentrate of the current invention includes alkanoyl N-methylglucamides of the formula




- where R^{10} is chosen from the group comprising C_6 and C_{22} alkyl
35 groups.

In another preferred embodiment the optional additives of the pearlescent concentrate of the current invention includes ether carboxylic acids of the formula



- 5 where R^{11} is chosen from the group includes C_6 to C_{22} alkyl groups
and m is from 1 to 120


- In a further preferred embodiment the optional additives of the pearlescent concentrate of the current invention
10 comprise ethoxylated sorbitan fatty acid esters of the formula



15

- where R^{12} is chosen from the group comprising
 C_6 to C_{22} alkyl groups
and w, v, u, t are from 4 to 20

- In a further preferred embodiment the optional additives
20 of the pearlescent concentrate of the current invention
comprise alpha-olefin sulphonates of formula



- where R^{13} is chosen from the group comprising
 C_6 to C_{22} alkyl groups
25 and M^2 is chosen from the group comprising Na & K.

Another optional additive to the pearlescent based concentrates of the current invention is between 0 and 2% of at least one inorganic salt including alkali metal halide salt such as NaCl, KCl, KI and the like.

- 30 The pearlescent based concentrates of the current invention may also be diluted with between 10 and 90% of the weight of the composition of water or other suitable solvent.

The pearlescent based concentrates of the current invention may also have added optional agents including perfumes, pigments, UV absorbers and antioxidants.

There is further provided a method of preparation of the
5 pearlescent based concentrates of the current invention comprising the steps of;

- (a) heating the components to a temperature above the highest melting point of the components but less than the boiling point of any solvents present and stirring
10 to form a homogeneous solution, and
- (b) cooling the stirring solution to a temperature of between 35°C and 45°C at a maximum cooling rate of 15°C/hour.

The pearlescent based concentrates of the current
15 invention may also be prepared by a method comprising the following steps;

- (a) heating the components to a temperature above the highest melting point of the components but less than the boiling point of any solvents present while stirring
20 vigorously to form a homogeneous solution,
- (b) rapidly cooling the stirring solution to between 60°C and 70°C, and
- (c) cooling the stirring solution to a temperature of between 35°C and 45°C at a maximum cooling rate of
25 15°C/hour.

The current invention will now be described with reference to the following non-limiting examples;

EXAMPLE 1

A pearlescent based concentrate of the following composition was prepared by the method described below.

<u>Component</u>	<u>Composition</u> (% w/w)
5 Decylpolysaccharide (50% a.i.)	36.00
Sodium lauryl sulphate (30% a.i.)	10.00
Ethylene Glycol mono-stearate	16.25
Sodium chloride	1.00
Water	36.75

10 Method

Ethylene glycol monostearate, decylpolysaccharide, alkyl sulphate, water and sodium chloride were heated to 80°C with vigorous stirring. The solution was maintained at 80°C with stirring for further 30 minutes before being rapidly cooled to 65°C. Once the solution temperature reached 65°C the stirring rate was slowed and the solution cooled at a controlled rate of 10°C per hour until the temperature reached 40°C. The stirrer was then stopped and the pearlescent blend allowed to cool to room temperature.

20 EXAMPLE 2

A pearlescent based concentrate of the following composition was prepared by the method described below.

<u>Component</u>	<u>Composition</u> (% w/w)
25 Dodecylpolysaccharide (50% a.i.)	25.00
Sodium lauryl sulphate (30% a.i.)	31.00
TERIC 12A23	3.00
Ethylene glycol mono-stearate	12.50
Sodium chloride	1.00
Water	27.50

30 Method

Ethylene glycol mono-stearate, dodecylpolysaccharide, sodium lauryl sulphate, TERIC 12A23 water and sodium chloride were heated to 80°C with vigorous stirring. Once the solution had reached 80°C, the stirring rate was slowed and the solution cooled at a controlled rate of 10°C per hour until the solution temperature had dropped to 40°C. When the solution temperature had reached 40°C the stirrer was stopped

and the pearlescent based composition was allowed to cool to room temperature. (TERIC is a registered trade mark of ICI Australia Operations Proprietary Limited).

EXAMPLE 3

5	<u>Component</u>	<u>Composition</u> (%w/w)
	Decylpolysaccharide (50% a.i.)	37.00
	Sodium lauryl sulphate	10.00
	Cetyl Stearyl alcohol	5.00
	Ethylene glycol monostearate	16.25
10	Water	32.75

Method

Ethylene glycol monostearate, decylpolysaccharide, sodium lauryl sulphate, cetyl-stearyl alcohol and water were heated to 75°C with vigorous stirring. The solution was cooled until the temperature reached 70°C at which point the rate of cooling was controlled at 15°C/hour and the stirring rate reduced. When temperature reached 40°C, stirring was stopped and the pearlescent based composition allowed to cool to room temperature.

20 EXAMPLE 4

	<u>Component</u>	<u>Composition</u> (%w/w)
	Dodecylpolysaccharide (50% a.i.)	36.00
	Ammonium lauryl sulphate (25% a.i.)	10.00
25	Disodium alkyl ethoxy sulphosuccinate (30% a.i.)	13.3
	Sodium chloride	1.0
	Ethylene glycol distearate	16.25
	Water	23.45

Method

30 Dodecylpolysaccharide, ammonium lauryl sulphate, disodium alkylethoxy sulphosuccinate and ethylene glycoldistearate were blended together to form a homogeneous mixture. Sodium chloride was then dissolved in the water and added to the mixture which was heated to 70°C with vigorous stirring. When the solution temperature reached 70°C the stirring rate was reduced and the solution cooled at a rate of 8°C/hr. When the solution temperature reached 40°C the

stirring was stopped and the resultant pearlescent based concentrate left to cool to room temperature.

EXAMPLE 5

<u>Component</u>	<u>Composition (%w/w)</u>
5 Decyl Polysaccharide	25.00
Sodium lauryl sulphate	20.00
Ethylene glycol monostearate	7.50
Sodium Chloride	1.00
Water	46.5

10 Method

The sodium chloride was dissolved in a portion of the water. The remaining components were blended together and the salt solution mixed in. The resultant mixture was heated to 70°C with rapid stirring then the stirring rate was reduced and the mixture cooled at a rate of 8°C per hour. When the mixture reached 40°C all stirring was ceased.

EXAMPLE 6

<u>Component</u>	<u>Composition (%w/w)</u>
Decyl Polysaccharide	25.00
20 Sodium lauryl sulphate	20.00
Sorbitan monolaurate	5.00
Ethylene glycol monostearate	15.00
Sodium Chloride	1.00
Water	34.00

25 Method

The above components were combined according to the method of Example 5 to form a pearlescent based concentrate.

EXAMPLE 7

<u>Component</u>	<u>Composition (%w/w)</u>
30 C8-10 Polysaccharide	5.00
Decyl Polysaccharide	25.00
Sodium lauryl sulphate	20.00
Ethylene glycol monostearate	15.00
Sodium Chloride	1.00
35 Water	34.00

Method

The above components were combined according to the method of Example 5 to form a pearlescent based concentrate.

EXAMPLE 8

5	<u>Component</u>	<u>Composition</u> (%w/w)
	Decyl Polysaccharide	30.00
	Ammonium lauryl sulphate	20.00
	Ethoxylated sorbitan monooleate	2.50
	Sodium Chloride	1.00
10	Ethylene glycol monostearate	15.00
	Water	31.50

Method

The above components were combined according to the method of Example 5 to form a pearlescent based concentrate.

15 EXAMPLE 9

	<u>Component</u>	<u>Composition</u> (%w/w)
	Decyl Polysaccharide	30.00
	Sodium lauryl sulphate	15.00
	Stearic acid	15.00
20	Sodium Chloride	1.00
	Water	39.00

Method

The above components were combined according to the method of Example 5 to form a pearlescent based concentrate.

25 EXAMPLE 10

	<u>Component</u>	<u>Composition</u> (%w/w)
	C16-18 Polysaccharide	10.00
	Decyl Polysaccharide	15.00
	Sodium lauryl sulphate	15.00
30	Ethylene glycol monostearate	15.00
	Sodium Chloride	1.00
	Water	44.00

Method


The above components were combined according to the
35 method of Example 5 to form a pearlescent based concentrate.

The pearlescent based concentrates of each example exhibited small crystals of relatively uniform size which give a lustrous appearance to the concentrate and good reflectance.

- 5 While the invention had been explained in relation to its preferred embodiments it is to be understood that various modifications thereof will become apparent to those skilled in the art upon reading the specification. Therefore, it is to be understood that the invention disclosed herein is
- 10 intended to cover such modifications as fall within the scope of the appended claims.

The claims defining the inventions are as follows:

1. A pearlescent based concentrate comprising:
between 5 and 40% by weight of the composition of
alkylpolysaccharide of formula




where R^1 is selected from the group comprising linear or
branched C_6 to C_{22} alkyl or alkenyl group

where G is selected from the group comprising a C_5 or C_6
saccharide and

10 x is from 1 to 10;

between 3 and 30% by weight of the composition of alkyl
sulphate and/or alkyl ether sulphate;

and between 5 and 20% by weight of pearlescing compound of
formula



where R^2 is chosen from the group including C_{16} to C_{22} fatty
acyl groups,

R^3 is chosen from the group comprising H or R^2 ,

n is 2 or 3 and

20 x is from 0 to 120.

2. A pearlescent based concentrate according to claim 1
comprising:

between 5 and 40% by weight of the composition of
alkylpolysaccharide of formula



where R^1 is selected from the group comprising linear or
branched C_6 to C_{22} alkyl or alkenyl group

where G is selected from the group comprising a C_5 or C_6
saccharide and

30 x is from 1 to 10;


between 3 and 30% by weight of the composition of alkyl
sulphate and/or alkyl ether sulphate;

and between 5 and 20% by weight of pearlescing acid of formula



where R is chosen from the group including C_{16} to C_{22} alkyl groups,


3. A pearlescent based concentrate comprising:
between 5 and 40% by weight of the composition of
alkylpolysaccharide of formula



10 where R^1 is selected from the group comprising linear or
branched C_6 to C_{22} alkyl or alkenyl group
where G is selected from the group comprising a C_5 or C_6
saccharide and

x is from 1 to 10;

15 between 3 and 30% by weight of the composition of alkyl
sulphate and/or alkyl ether sulphate;
and between 5 and 20% by weight of pearlescing esters of
formula




20 where R^2 is chosen from the group including C_{16} to C_{22} fatty
acyl groups,

R^3 is chosen from the group comprising H or R^2 ,

n is 2 or 3 and

x is from 1 to 120.

25 4. A pearlescent based concentrate according to claim 3
which further includes secondary alkane sulphonates of the
formula




where R^4 is chosen from the group comprising

30 C_6 to C_{22} alkyl groups

and M is chosen from the group comprising alkali salts.

5. A pearlescent based concentrates according to claim 3
wherein said fatty alcohol ethoxylate is of formula




35 where R^5 is chosen from the group including

C_6 and C_{22} alkyl or alkenyl groups

and y is from 1 to 120.

6. A pearlescent based concentrate according to claim 3 which further includes between 0 and 30% by weight of the composition of one or more optional additives chosen from the group including:

alkylsulphosuccinate or polyoxyalkylenealkylsulphosuccinate of the formula




- where R^6 is chosen from the group including C_8 to C_{20} linear or branched alkyl groups,

R^7 is H or CH_3 ,

- M and M^1 are independently chosen from the group comprising alkali metals, alkaline earth metals, NH_4^+ , C_1 to C_3 alkyl groups, substituted ammonia and hydroxy substituted C_2 or C_3 alkyl substituted ammonia and

n is from 0 to 8;

sorbitan fatty acid esters of formula,



- where R^8 is chosen from the group including C_{12} to C_{18} alkyl group;

secondary alkane sulphonates; fatty alcohol alkoxyates; fatty acid alkoxyates; fatty alcohols; alkanoyl N-methylglucamides; ether carboxylic acids; ethoxylated sorbitan fatty acid esters; and alpha-olefin sulphonates.

7. A pearlescent based concentrate according to claim 3 which further includes fatty alcohols of the formula



where R^9 is chosen from the group comprising C_6 to C_{22} alkyl groups.


8. A pearlescent based concentrate according to claim 3 which further includes alkanoyl N-methylglucamides of the formula



- $CH_3-N-CH_2(CHOH)_4CH_2OH$


where R^{10} is chosen from the group consisting C_6 and C_{22} alkyl groups.

9. A pearlescent based concentrate according to claim 3 which further includes ether carboxylic acids of the formula



where R^{11} is chosen from the group including C_6 to C_{22} alkyl groups and m is from 1 to 120.


10. A pearlescent based concentrate according to claim 3 which further includes ethoxylated sorbitan fatty acid esters of the formula



where R^{12} is chosen from the group comprising C_6 to C_{22} alkyl groups

and w, v, u, t are from 4 to 20

11. A pearlescent based concentrate according to claim 3 which further includes alpha-olefin sulphonates of formula



where R^{13} is chosen from the group comprising

C_6 to C_{22} alkyl groups

and M^2 is chosen from the group comprising Na & K.


12. A pearlescent based concentrate according to claim 3 comprising between 0 and 2% of the weight of the composition of inorganic salt.

13. A pearlescent based concentrate according to claim 12 wherein said inorganic salt is an alkali metal halide salt.

14. A pearlescent based concentrate according to claim 3 comprising between 10 and 90% of the weight of the composition of solvent.

15. A pearlescent composition comprising a pearlescent based concentrate according to any of the previous claims and at least one solvent in the ratio of between 10:90 and 90:10 weight %.

- 5 16. A pearlescent based concentrate according to claim 1 or 2 which further includes secondary alkane sulphonates of the formula




where R^4 is chosen from the group comprising

- 10 C_6 to C_{22} alkyl groups

and M is chosen from the group comprising alkali salts.

17. A pearlescent based concentrates according to claim 1 or 2 wherein said fatty alcohol ethoxylate is of formula




- 15 where R^5 is chosen from the group including

C_6 and C_{22} alkyl or alkenyl groups

and y is from 1 to 120.

18. A pearlescent based concentrate according to claim 1 or 2 which further includes between 0 and 30% by weight of the composition of one or more optional additives chosen from the group including:

alkylsulphosuccinate or polyoxyalkylenealkylsulphosuccinate of the formula



- 25 where R^6 is chosen from the group including C_8 to C_{20} linear or branched alkyl groups,


R^7 is H or CH_3 ,

M and M^1 are independently chosen from the group comprising alkali metals, alkaline earth metals, NH_4^+ , C_1 to C_3 alkyl

- 30 groups, substituted ammonia and hydroxy substituted C_2 or C_3 alkyl substituted ammonia and

n is from 0 to 8;

sorbitan fatty acid esters of formula,



- 35 where R^8 is chosen from the group including C_{12} to C_{18} alkyl group;

secondary alkane sulphonates; fatty alcohol alkoxyates; fatty acid alkoxyates; fatty alcohols; alkanoyl N-methylglucamides; ether carboxylic acids; ethoxylated sorbitan fatty acid esters; and alpha-olefin sulphonates.

- 5 19. A pearlescent based concentrate according to claim 1 or 2 which further includes fatty alcohols of the formula



where R^9 is chosen from the group comprising C_6 to C_{22} alkyl groups.


- 10 20. A pearlescent based concentrate according to claim 1 or 2 which further includes alkanoyl N-methylglucamides of the formula



- 15 $CH_3-N-CH_2(CHOH)_4CH_2OH$


where R^{10} is chosen from the group consisting C_6 and C_{22} alkyl groups.

21. A pearlescent based concentrate according to claim 1 or 2 which further includes ether carboxylic acids of the
20 formula



where R^{11} is chosen from the group including C_6 to C_{22} alkyl groups and m is from 1 to 120.


22. A pearlescent based concentrate according to claim 1 or
25 2 which further includes ethoxylated sorbitan fatty acid esters of the formula



30

where R^{12} is chosen from the group comprising C_6 to C_{22} alkyl groups and w, v, u, t are from 4 to 20

23. A pearlescent based concentrate according to claim 3 which further includes alpha-olefin sulphonates of formula



where R^{13} is chosen from the group comprising

5 C_6 to C_{22} alkyl groups

and M^2 is chosen from the group comprising Na & K.

24. A method of preparing a pearlescent based concentrates of any of claims 3 to 15 comprising the steps of;

10 (a) heating the components to a temperature above the highest melting point of the components but less than the boiling point of any solvents present and stirring to form a homogeneous solution, and

(b) cooling the stirring solution to a temperature of between 35°C and 45°C at a maximum cooling rate of 15°C/hour.

25. A method of preparing a pearlescent based concentrates of any of claims 3 to 15 comprising the steps of;

20 (a) heating the components to a temperature above the highest melting point of the components but less than the boiling point of any solvents present while stirring vigorously to form a homogeneous solution,

(b) rapidly cooling the stirring solution to between 60°C and 70°C, and

25 (c) cooling the stirring solution to a temperature of between 35°C and 45°C at a maximum cooling rate of 15°C/hour.

26. A method of preparing a pearlescent based concentrates of any of claims 1, 2 or 16 to 23 comprising the steps of;

30 (a) heating the components to a temperature above the highest melting point of the components but less than the boiling point of any solvents present and stirring to form a homogeneous solution, and

(b) cooling the stirring solution to a temperature of between 35°C and 45°C at a maximum cooling rate of 15°C/hour.

27. A method of preparing a pearlescent based concentrates of any of claims 1, 2 or 16 to 23 comprising the steps of;
- 5 (a) heating the components to a temperature above the highest melting point of the components but less than the boiling point of any solvents present while stirring vigorously to form a homogeneous solution,
- (b) rapidly cooling the stirring solution to between 60°C and 70°C, and
- 10 (c) cooling the stirring solution to a temperature of between 35°C and 45°C at a maximum cooling rate of 15°C/hour.
28. A pearlescent based concentrate substantially as herein described with reference to the examples.
29. A method of preparing a pearlescent based concentrate as
15 herein described with reference to the examples.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU 94/00421

A. CLASSIFICATION OF SUBJECT MATTERInt. Cl.⁵ A61K 7/50, 7/075, C11D 17/00, 9/26, 9/44

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC A61K 7/50, 7/075, 7/06, 7/07, C11D 9/26, 9/44.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
AU : IPC as above

Electronic data base consulted during the international search (name of data base, and where practicable, search terms used)

DERWENT : } (Pearl:) (Soap)
JAPIO : } (or) and (or Shampoo)
CASM : } (Pearlesc:) (or Hair Conditioner)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
Y	WO,A, 86/05390 (HELENE CURTIS INDUSTRIES, INC.) 25 September 1986 (25.09.86) Page 5 lines 30-35, Page 18 lines 34-Page 19 line 27 and page 23 Lines 9 to 34	1-29
P, Y	WO,A, 93/15171 (DEUTSCHE ICI GMBH et al.) 5 August 1993 (05.08.93) Page 2 line 4 - page 3 line 18, Page 6 lines 16-32	1-29
Y	US,A, 4438096 (HELENE CURTIS INDUSTRIES, INC.) 20 March 1984 (20.03.84) Page 3 lines 24-62, Page 3 4 lines 9-17	1-29

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.*** Special categories of cited documents :**

"A" document defining the general state of the art which is not considered to be of particular relevance
"E" earlier document but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"&" document member of the same patent family

Date of the actual completion of the international search
23 September 1994 (23.09.94)

Date of mailing of the international search report


13 Oct 1994 (13.10.94)

Name and mailing address of the ISA/AU

AUSTRALIAN INDUSTRIAL PROPERTY ORGANISATION
PO BOX 200
WODEN ACT 2606
AUSTRALIA

Facsimile No. 06 2853929

Authorized officer:



JOHN G. HANSON

Telephone No. (06) 2832262

INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU 94/00421

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate of the relevant passages	Relevant to Claim No.
Y	US,A, 5017305 (HENKEL KOMMANDITGESELLSCHAFT AUF AKTIEN) 21 May 1991 (21.05.91)	1-29
A	AU,A, 26058/92 (KAO CORPORATION) 8 April 1993 (08.04.93)	
A	EP,A, 323594 (KAO CORPORATION) 12 July 1989 (12.07.89)	
A	EP,A, 512744 (UNILEVER PLC) 11 November 1992 (11.11.92)	

PCT/AU 94/00421

Patent Document Cited in Search Report		Patent Family Member					
WO	8605390	AU	55126/86	AU	594044	DK	5403/86
		EP	215025	JP	62501192	US	4654207
WO	9315171	AU	11813/92				
US	5017305	EP	268992	JP	63150214	US	4824594
AU	26058/92	EP	535693	JP	5117687		
EP	323594	JP	1176445	US	4959206		
EP	512744	GB	9109693	JP	5139940		
END OF ANNEX							

